

# Particle acceleration by magnetized rotating black hole

**Vladimír Karas**

Astronomical Institute, Czech Academy of Sciences




---

**07-11 SEPTEMBER 2026 | GARCHING BEI MÜNCHEN**


---

*Annual Meeting of the German Astronomical Society*



Motivation: Energy spectrum of primary cosmic rays

# Motivation: magnetic reconnection in ergosphere?



# The model: rotating black hole

Kerr metric describing the geometry of the spacetime around the rotating black hole is expressed in Boyer-Lindquist coordinates  $x^\mu = (t, r, \theta, \varphi)$  as follows (Misner et al. 1973):

$$ds^2 = -\frac{\Delta}{\Sigma} [dt - a \sin \theta d\varphi]^2 + \frac{\sin^2 \theta}{\Sigma} [(r^2 + a^2)d\varphi - a dt]^2 + \frac{\Sigma}{\Delta} dr^2 + \Sigma d\theta^2, \quad (1)$$

where

$$\Delta \equiv r^2 - 2Mr + a^2, \quad \Sigma \equiv r^2 + a^2 \cos^2 \theta. \quad (2)$$

Coordinate singularity at  $\Delta = 0$  corresponds to outer/inner horizon of the black hole  $r_\pm = M \pm \sqrt{M^2 - a^2}$ . Rotation of the black hole is measured by the spin parameter  $a \in \langle -M, M \rangle$ . Here we only consider  $a \geq 0$  without the loss of generality.

We employ the test-field solution of Maxwell's equations for a weakly magnetized Kerr black hole immersed in an asymptotically uniform magnetic field specified by the component  $B_z$  parallel to the spin axis and the perpendicular component  $B_x$ . The electromagnetic vector potential  $A_\mu$  is given as follows (Bičák & Janiš 1985):

$$A_t = \frac{B_z a M r}{\Sigma} (1 + \cos^2 \theta) - B_z a + \frac{B_x a M \sin \theta \cos \theta}{\Sigma} (r \cos \psi - a \sin \psi), \quad (3)$$

$$A_r = -B_x (r - M) \cos \theta \sin \theta \sin \psi, \quad (4)$$

$$A_\theta = -B_x a (r \sin^2 \theta + M \cos^2 \theta) \cos \psi - B_x (r^2 \cos^2 \theta - M r \cos 2\theta + a^2 \cos 2\theta) \sin \psi, \quad (5)$$

$$A_\varphi = B_z \sin^2 \theta \left[ \frac{1}{2} (r^2 + a^2) - \frac{a^2 M r}{\Sigma} (1 + \cos^2 \theta) \right] - B_x \sin \theta \cos \theta \left[ \Delta \cos \psi + \frac{(r^2 + a^2) M}{\Sigma} (r \cos \psi - a \sin \psi) \right], \quad (6)$$

where  $\psi$  denotes the azimuthal coordinate of Kerr ingoing coordinates, which is expressed in Boyer–Lindquist coordinates as follows:

$$\psi = \varphi + \frac{a}{r_+ - r_-} \ln \frac{r - r_+}{r - r_-}. \quad (7)$$

# Conditions for the escape of charged particles

Effective potential expressing the minimal allowed energy of charged test particles in a non-axisymmetric magnetosphere of a rotating black hole may be derived in the rest frame of a static observer (Kopáček & Karas 2018c). Tetrad vectors of this frame are given as (Semerák 1993):

$$e_{(t)}^\mu = \left[ \frac{\Sigma^{1/2}}{\chi}, 0, 0, 0 \right], \quad e_{(r)}^\mu = \left[ 0, \frac{\Delta^{1/2}}{\Sigma^{1/2}}, 0, 0 \right], \quad (10)$$

$$e_{(\theta)}^\mu = \left[ 0, 0, \frac{1}{\Sigma^{1/2}}, 0 \right], \quad e_{(\varphi)}^\mu = \frac{\chi}{\sin \theta \Delta^{1/2} \Sigma^{1/2}} \left[ \frac{-2aMr \sin^2 \theta}{\chi^2}, 0, 0, 1 \right], \quad (11)$$

where  $\chi^2 \equiv \Delta - a^2 \sin^2 \theta$ .

The static frame is employed to express the effective potential:

$$V_{\text{eff}}(r, \theta, \varphi) = \left( -\beta + \sqrt{\beta^2 - 4\alpha\gamma} \right) / 2\alpha, \quad (12)$$

with the coefficients defined as:

$$\alpha = [e_{(t)}^t]^2, \quad \beta = 2qA_t e_{(t)}^t, \quad \gamma = q^2 [e_{(t)}^t]^2 A_t^2 - 1. \quad (13)$$


# Escape in oblique configuration

Once the charge is introduced, the value of effective potential (12) changes accordingly. In order to study trajectories of escaping particles, we examine the behavior of the potential for  $r \gg M$ . In particular, for the initially neutral particle with energy  $E_{\text{Kep}}$  ionized in the equatorial plane at  $r_0$  we obtain the following relation valid in the asymptotic region:

$$E - V_{\text{eff}}|_{r \gg M} = E_{\text{Kep}} - 1 - \frac{qB_z a}{r_0} + \mathcal{O}(r^{-1}). \quad (18)$$


Motion is possible only for  $E \geq V_{\text{eff}}$ . Since  $E_{\text{Kep}} < 1$  with finite  $r_0$  and  $a \geq 0$  is considered, we observe that (i) particles may only escape for  $qB_z < 0$ , (ii) the escape is possible only for  $a \neq 0$ , (iii) asymptotic velocity of escaping particles is an increasing function of parameters  $|qB_z|$  and  $a$ , and a decreasing function of  $r_0$ , and (iv) the escape is not allowed for the perpendicular inclination,  $\alpha \equiv \arctan(B_x/B_z) = \pi/2$ .

# Rotating black hole, oblique magnetic field




Effect of misalignment.

# Magnetic null points




Field lines in equatorial plane

Karas et al. (CQG, 2009, 2012)




**Figure 1.** Different types of the trajectories launched from the equatorial plane  $(x, y)$  are plotted with respect to the magnetic inclination angle  $\alpha$ . Color-coding: blue for plunging orbits, red for stable ones (bound to the black hole) and yellow for escaping trajectories. Green circle denotes the ISCO. Inner black region marks the horizon of the black hole. Parameters of the system are  $a = 0.98$  and  $qB = -5$ . Magnetic field is inclined in the positive  $x$ -direction.




**Figure 6.** The structure of the escape zone ( $qB = -4.1$ ,  $\alpha = 14^\circ$  and  $\varphi_0 = 0$ ) in the relevant range of the spin parameter  $a$  and initial radius  $r_0$  is explored. Going from the top-left to the bottom-right panel, the portions of the plots (marked by black rectangles) are magnified progressively in the subsequent panels revealing the complex structure. Green line represents the ISCO, black color shows the horizon of the black hole. Color-coding of trajectories as in Figs. 1-3.

To explore the effects of the black hole charge, we begin with the aligned (Wald) field.




**Figure 1.** Evolution of escape zones of charged particles affected by the Coulomb repulsion of charged black hole. Color-coding of trajectories: blue for plunging orbits, red for stable ones (bound to the black hole) and yellow for escaping trajectories. The inner black region marks the horizon of the black hole. The green line denotes the ISCO of neutral particles, while the cyan line marks MBSO. Parameters of the system are  $qB = -1$  and inclination  $\alpha = 0$  while  $qQ$  varies as indicated.


See also Kopáček & Karas (2026), to be submitted




**Figure 3.** Maximum Lorentz factor achieved within the escape zones as a function of charge product  $qQ$  compared with a theoretical limit for given values of parameters, as determined by eq. (21) (dashed lines).



**Figure 4.** Values of initial radial coordinate of maximally accelerated escaping particle within escape zone as a function of charge product  $qQ$ .



**Figure 10.** Maximal value of the final Lorentz factor of escaping particles from the vicinity of the Kerr black hole with Wald charge as a function of parameter  $qB$ . Numerically obtained actual values detected within escape zones (black) vs. estimated theoretical maximum (red).



**Figure 5.** The Innermost Escape Zone affected by the inclination of the magnetic field. Color-coding of trajectories is the same as in Fig. 1. Parameters of the system are  $qB = -1$ ,  $qQ = 1.2$ ,  $\varphi(0) = 0$ , and inclination angle  $\alpha$  varies as indicated.

# Summary

We have computed the final Lorentz factor  $\gamma$  of escaping particles confirming that the highest  $\gamma$  is achieved in the innermost region of the primary escape zone (with the lowest allowed  $r_0$ ). For a particular (realistically small;  $\alpha \approx 6^\circ$ ) value of inclination and fixed value of spin ( $a = 0.98$ ), we searched for the maximal  $\gamma$ . Increasing the value of magnetization up to  $|qB| = 10^3$  we confirmed that (unlike axisymmetric configuration) ultrarelativistic velocities with  $\gamma \gg 1$  may be achieved. While the acceleration of the particle is actually powered by the parallel component  $B_z$ , the perpendicular component  $B_x$  acts as an extra perturbation which considerably increases the probability of sending the particles on escaping trajectories and allows the outflow also in cases which are excluded in the aligned setup.

- Kopáček & Karas, *The Astrophysical Journal*, 966, 226 (2024)
- -----"-----, *Proc. IAU Symp. #405 (Brno, CZ)*, in press (2026)
- -----"-----, *The Astrophysical Journal*, in prep. (2026)